



Restoring Sockeye Salmon to the Yakima River Basin.

Mark V. Johnston¹, David E. Fast¹, Brian Saluskin¹, William J. Bosch¹, and Stephen J. Grabowski².

¹Yakama Nation – Yakima/Klickitat Fisheries Project, Toppenish, WA. ²U.S. Bureau of Reclamation, Boise, ID.



Abstract

Returns of sockeye salmon to the upper Columbia Basin have numbered 50,000 or fewer in 14 of the past 22 years. Dam counts indicate that sockeye are declining by an average of 830 fish per year. Of the historic sockeye nursery lake habitat in the Upper Columbia, only about 4% is presently utilized with only two (Wenatchee and Osoyoos) of 12 historic nursery lakes presently producing fish. Four nursery lakes in the Yakima River Basin, which historically produced an estimated annual return of about 200,000 sockeye, were removed from production in the early 1900s when irrigation storage dams were constructed without passage. Work conducted by the National Marine Fisheries Service from 1987 to 1993 in Lake Cle Elum returned from 4 to 20 sockeye adults to the base of Cle Elum Dam demonstrating that sockeye restoration was feasible with sufficient passage modifications. In 1994 the Yakima River Basin Water Enhancement Project Act was passed providing for increased storage capacity in Cle Elum reservoir including provisions for developing fish passage alternatives. The Yakama Nation is now working with the Bureau of Reclamation to conduct additional feasibility work using a temporary juvenile passage flume and coho salmon as surrogates. Over 25% of the smolts released into Lake Cle Elum in 2006-2007 successfully migrated using the flume and 1.5% of the known 2006 outmigrants returned as adults to Prosser Dam in 2007. Based on the results of this prior work, the Yakama Nation is working to resolve logistical issues so that sockeye fry plants can begin in Lake Cle Elum in 2009.

In a January 2007 memorandum discussing the role of large extirpated areas in recovery, the Interior Columbia Technical Recovery Team noted that "Snake River sockeye are currently restricted to a single extant population. The probability of long-term persistence of the [evolutionarily significant unit] ESU will be greatly enhanced with additional populations. In fact, the ESU cannot meet the minimum ESU biological viability criteria established by the TRT without multiple viable populations."

The Subbasin Plan for the Upper Middle Columbia River Mainstem cited the Northwest Power and Conservation Council's (NPCC) 2000 Fish and Wildlife program in establishing the vision for that subbasin: "There is an obligation to provide fish and wildlife mitigation where habitat has been permanently lost because of hydroelectric development. Restoration of anadromous fish into areas blocked by dams should be actively pursued where feasible." Clearly, restoring sockeye to additional Columbia River nursery lakes is consistent with regional mitigation obligations and the best available science on Columbia River salmon recovery.

Restoration Potential in the Yakima Subbasin




Fig 2. Location of Historic Sockeye Nursery Lakes in the Yakima River Basin.

Four nursery lakes in the Yakima River Basin (Fig. 2), which historically produced an estimated annual return of about 200,000 sockeye, were removed from production in the early 1900s when irrigation storage dams were constructed without passage.

Between 1987-1993, the National Marine Fisheries Service (NMFS), Northwest Fisheries Science Center, annually outplanted about 100-350,000 sockeye salmon juveniles at Cle Elum Lake to study the feasibility of restoring sockeye salmon to the Yakima River Basin (Flagg et al. 2000). Merwin trapping efforts in Lake Cle Elum during spring 1990-1993 indicated that in most years juvenile sockeye salmon outplanted to the lake survived and grew well. Investigation of fish passage from the tailrace of Cle Elum Dam through the Yakima River system during 1989-1993 suggested there were no severe blockages to migration of sockeye salmon in the Yakima River system downstream of Cle Elum Dam during the normal late-March to early-May period of smolt outmigration for sockeye salmon in the Columbia River Basin. However, fish passage studies at Cle Elum Dam during spring 1989-1993 suggested a fish passage problem at the dam during much of the outmigration period. The results of the studies suggested that, if mechanical traps and fishways were constructed at Cle Elum Dam, sockeye salmon could be restored to Cle Elum Lake.

Current Restoration Effort

As part of water storage improvements under Section 1206 of the 1994 Yakima River Basin Water Enhancement Project Act, Title XII of Public Law 103-434, the Yakama Nation, with the cooperation of the U.S. Bureau of Reclamation, is now actively pursuing the restoration of anadromous fish passage above Cle Elum Dam. The BOR estimated sockeye smolt production potential of 400,000 to 1.6 million fish in the Cle Elum Lake watershed (Fig. 3), with a projected return of 30,000 to 50,000 adult spawners assuming average survival and median pool elevation (Grabowski 2007). Based on this and other feasibility work, a temporary juvenile fish passage flume (Fig. 4) was constructed at Cle Elum Dam in 2006. The Yakama Nation tested passage through this flume in 2006-07 using coho salmon (*O. kisutch*) as surrogates. Approximately 10,000 coho smolts were released annually and passive integrated transponder (PIT) tags were used to evaluate survival. Over 25% of the smolts released into Lake Cle Elum in 2006-2007 successfully migrated using the flume and 1.5% of the known 2006 outmigrants returned as adults to Prosser Dam in 2007.




Fig 3 (L). Lake Cle Elum Watershed and Fig. 4 (above) Temporary Passage Flume.

Based on the results of this feasibility work, the Yakama Nation is working with relevant co-managers to implement a plan to collect up to 300 pairs of sockeye (Osoyoos population, Okanogan R.) at Wells Dam in 2008, spawn and rear the progeny of these fish at a facility in the Upper Columbia Basin, and plant these fry in Lake Cle Elum in 2009 when they have passed all necessary fish health inspections. Lake Osoyoos sockeye are believed to be the best suited donor population as environmental conditions in their migration corridor are considered most similar to present conditions in the Yakima Basin during the adult sockeye migration period. Adult returns from these releases would be collected at Roza Dam, transported and released above Cle Elum Dam (short-term) or used as broodstock once an integrated supplementation program is developed (long-term). The Yakama Nation will continue to work with the BOR to ensure that permanent juvenile and adult passage facilities are constructed at Cle Elum Dam so that this restoration effort will eventually result in a self-sustaining population.

References

Flagg, T. A., T. E. Ruehle, L. W. Harrell, J. L. Mighell, C. R. Pasley, A. J. Novotny, E. Stettin, C. W. Sims, D. B. Dey, and C. V. W. Mahlink - National Marine Fisheries Service, Seattle, WA. 2000. Cle Elum Lake Anadromous Salmon Restoration Feasibility Study. Summary of Research, 2000 Final Report to Bonneville Power Administration, Portland, OR, Contract No. 86A164840, Project No. 86-045, 118 electronic pages (BPA Report DOE/BP-64840-4).

Grabowski, S. 2007. Assessment of Sockeye Salmon Production Potential in the Cle Elum River Basin, Storage Dam Fish Passage Study, Yakama Project, Washington. Technical Series No. PN-YDFP-008, Bureau of Reclamation, Boise, Idaho. March 2007.

ICTRT memo re role of large extirpated areas in recovery: http://www.nwrc.gov/ictrt/documents/attach1_viability_3_19_07.pdf

Independent Scientific Review Panel (ISRP). 2006. FY 2007-09 Final Review of Proposals. Council document ISRP 2006-6. Northwest Power and Conservation Council. Portland, OR. Available on-line at: <http://www.nwrcouncil.org/issrp/issrp06-6.htm>

Upper Middle Columbia River Mainstem Subbasin Plan Executive Summary, available on-line at: http://www.nwrcouncil.org/fw/subbasinplanning/uppermiddlecolumbia/plan/intro/overview_screen.pdf

WDFW (Washington Department of Fish and Wildlife)/ODFW (Oregon Department of Fish and Wildlife). 2002. Status Report, Columbia River Fish Runs and Fisheries, 1938-2000. The Joint Columbia River Management Staffs of Washington Department of Fish and Wildlife and Oregon Department of Fish and Wildlife. August 2002. Vancouver, WA. Available on-line at: http://wdfw.wa.gov/fish/columbia/2000_status_report_text.pdf

Columbia R. Sockeye in Decline




Fig 1. Adult Counts of Sockeye at Bonneville and McNary or Priest Rapids Dams, 1938 to present.

A simple trend analysis indicates that Sockeye (*Oncorhynchus nerka*) populations escaping to upper Columbia Basin spawning areas are declining by an average of 830 fish per year (Fig. 1). With the exception of a few years in the mid-1980s, Treaty harvest of sockeye in Zone 6 fisheries has been severely restricted since the mid-1970s (Fig. 1). Historically, the Columbia River sockeye run was as large as 3 million fish (WDFW/ODFW 2002). The recent 15-year average escapement at Priest Rapids Dam was fewer than 50,000 sockeye compared to a *U.S. v Oregon* escapement goal of 65,000. A large part of the decline is because Columbia and Snake River dams blocked access to historic nursery lake habitats, reducing available habitats by 96% (WDFW/ODFW 2002).

Need for Additional Populations and Restoration Efforts

Since the listing of Snake River sockeye salmon in 1991, the Columbia Basin's sockeye restoration efforts have consisted primarily of a captive broodstock program focused on restoring sockeye to Stanley Basin. In 2006, the Independent Scientific Review Panel (ISRP) concluded this effort was unlikely to succeed (ISRP 2006-6).